arXiv:1607.05104v1 [math.CA] 14 Jul 2016

Some Integral Inequalities For Functions Whoose Second Derivatives Are $\varphi-$Convex By Using Fractional Integrals

M. Esra YILDIRIM, Abdullah AKKURT, AND Hseyin YILDIRIM

Abstract. In this paper, we obtain new estimates on generalization of Hermite-Hadamard, Simpson and Ostrowski type inequalities for functions whose second derivatives is $\varphi-$convex via fractional integrals.

1. Introduction

The following inequality is called Hermite-Hadamard Inequality;

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a}\int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2}, \tag{1.1}$$

where $f : I \subseteq \mathbb{R} \to \mathbb{R}$ is a convex function and $a, b \in I$ with $a < b$. If f is concave, then both inequalities hold in the reversed direction .

The inequality 1.1 inequality was first discovered by Hermite in 1881 in the Journal Mathesis. This inequality was known as Hermite-Hadamard Inequlity, because this inequality was found by Mitrinovic Hermite and Hadamard' note in Mathesis in 1974.

The inequality 1.1 is studied by many authors, see ([1]-[7], [9]-[11], [13], [16]-[22]) where further references are listed.

Firstly, we need to recall some concepts of convexity concerning our work.

Definition 1. [6] *A function $f : I \subset \mathbb{R} \to \mathbb{R}$ is said to be convex on I if inequality*

$$f(ta + (1-t)b) \leq tf(a) + (1-t)f(b), \tag{1.2}$$

holds for all $a, b \in I$ and $t \in [0, 1]$.

Definition 2. [8] *Let $s \in (0, 1]$. A function $f : I \subseteq \mathbb{R}_0 = [0, \infty) \to \mathbb{R}$ is said to be $s-$convex in the second sense if*

$$f(ta + (1-t)b) \leq t^s f(a) + (1-t)^s f(b), \tag{1.3}$$

holds for all $a, b \in I$ and $t \in [0, 1]$.

Tunç and Yildirim in [22] introduced the following defination as follows:

Definition 3. *A function $f : I \subseteq \mathbb{R} \to \mathbb{R}$ is said to belong to the class of $MT(I)$ if it is nonnegative and for all $x, y \in I$ and $t \in (0, 1)$ satisfies the inequality;*

$$f(tx + (1-t)y) \leq \frac{\sqrt{t}}{2\sqrt{1-t}} f(x) + \frac{\sqrt{1-t}}{2\sqrt{t}} f(y).$$

1991 *Mathematics Subject Classification.* 26D15, 26A51, 26A33, 26D10.

Key words and phrases. Hermite–Hadamard inequality; Ostrowski inequality; Simpson inequality; Riemann–Liouville fractional integral, h -convex functions.

M.E. Yildirim was partially supported by the Scientific and Technological Research Council of Turkey (TUBITAK Programme 2228-B).

Dragomir in [3] introduced the following defination as follows:

Definition 4. [3] *Let $\varphi : (0,1) \to (0,\infty)$ be a measurable function. We say that the function $f : I \to [0,\infty)$ is a $\varphi-$convex function on the interval I if $x, y \in I$ we have*

$$f(tx + (1-t)y) \leq t\varphi(t) f(x) + (1-t)\varphi(1-t) f(y).$$

Remark 1. *According to defination 4 for the special choose of φ we can obtain following*

If we take $\varphi(t) \equiv 1$, we obtain classical convex.
If we take $\varphi(t) = t^{s-1}$, we obtain $s-$convex.
If we take $\varphi(t) = \dfrac{1}{2\sqrt{t}\sqrt{1-t}}$, we obtain $MT-$convex.

Now, we will give some definitions and notations of fractional calculus theory which are used later in this paper. Samko et al. in [15] used following definations as follows:

Definition 5. [15] *The Riemann-Liouville fractional integrals $J^{\alpha}_{a^+} f$ and $J^{\alpha}_{b^-} f$ of order $\alpha > 0$ with $a \geq 0$ are defined by*

$$J^{\alpha}_{a^+} f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t)dt, \ x > a \tag{1.4}$$

and

$$J^{\alpha}_{b^-} f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t)dt, \ x < b \tag{1.5}$$

where $f \in L_1[a,b]$, respectively. Note that, $\Gamma(\alpha)$ is the Gamma function and $J^0_{a^+} f(x) = J^{\alpha}_{b^-} f(x) = f(x)$.

Definition 6. [15] *The Euler Beta function is defined as follows:*

$$\beta(x,y) = \int_0^1 t^{x-1}(1-t)^{y-1} dt, \ x, y > 0.$$

The incomplate beta function is defined as follows:

$$\beta(a,x,y) = \int_0^a t^{x-1}(1-t)^{y-1} dt, \ x, y > 0, \ 0 < \alpha < 1.$$

2. Main results

Througout this paper, we use S_f as follows;

$$\begin{aligned} S_f(x,\lambda,\alpha;a,b) \ &\equiv (1-\lambda)\left\{\tfrac{(b-x)^{\alpha+1}-(x-a)^{\alpha+1}}{b-a}\right\} f'(x) \\ &+ (1+\alpha-\lambda)\left\{\tfrac{(x-a)^{\alpha}+(b-x)^{\alpha}}{b-a}\right\} f(x) \\ &+ \lambda\left\{\tfrac{(x-a)^{\alpha}(f(a)+(b-x)^{\alpha} f(b)}{b-a}\right\} \\ &- \tfrac{\Gamma(\alpha+2)}{b-a}\left\{J^{\alpha}_{x^-} f(a) + J^{\alpha}_{x^+} f(b)\right\}, \end{aligned}$$

for any $x \in [a,b]$, $\lambda \in [0,1]$ and $\alpha > 0$.

In ([14]), Jaekeun Park established the following lemma which is necessary to prove our main results:

Lemma 1. *Let $f: I \subseteq \mathbb{R} \to \mathbb{R}$ be a twice differentiable function on the interior I^0 of an interval I such that $f'' \in L_1[a,b]$, where $a,b \in I$ with $a < b$. Then, for any $x \in [a,b]$, $\lambda \in [0,1]$ and $\alpha > 0$ we have*

$$\begin{aligned} S_f(x,\lambda,\alpha;a,b) &= \tfrac{(x-a)^{\alpha+2}}{b-a}\int_0^1 t\left(\lambda - t^\alpha\right) f''\left(tx+(1-t)a\right)dt \\ &\quad + \tfrac{(b-x)^{\alpha+2}}{b-a}\int_0^1 t\left(\lambda - t^\alpha\right) f''\left(tx+(1-t)b\right)dt. \end{aligned}$$

Theorem 1. *Let $\varphi : (0,1) \to (0,\infty)$ be a measurable function. Assume also that $f : I \subset [0,\infty) \to \mathbb{R}$ be a twice differentiable function on the interior I^0 of an interval I such that $f'' \in L_1[a,b]$, where $a,b \in I^0$ with $a < b$. If $|f''|^q$ is $\varphi-$convex on $[a,b]$ for some fixed $q \geq 1$, then for any $x = ta + (1-t)b, t \in [0,1], \lambda \in [0,1]$, and $\alpha > 0$:*

$$\begin{aligned} |S_f(x,\lambda,\alpha,t,\varphi;a,b)| &\leq A_1^{1-\frac{1}{q}}(\alpha,\lambda)\left[\tfrac{(x-a)^{\alpha+2}}{b-a}\left\{A_2(\alpha,\lambda,t,\varphi)\left|f''(x)\right|^q\right.\right. \\ &\quad \left. + A_3(\alpha,\lambda,t,\varphi)\left|f''(a)\right|^q\right\}^{\frac{1}{q}} \qquad (2.1) \\ &\quad \left. + \tfrac{(b-x)^{\alpha+2}}{b-a}\left\{A_2(\alpha,\lambda,t,\varphi)\left|f''(x)\right|^q + A_3(\alpha,\lambda,t,\varphi)\left|f''(b)\right|^q\right\}^{\frac{1}{q}}\right], \end{aligned}$$

the above inequality for fractional integrals holds, where

$$\begin{aligned} A_1(\alpha,\lambda) &= \tfrac{\alpha\lambda^{1+\frac{2}{\alpha}}+1}{\alpha+2} - \tfrac{\lambda}{2}, \\ A_2(\alpha,\lambda,t,\varphi) &= \int_0^1 \left|t\left(\lambda - t^\alpha\right)\right| t\varphi(t)\,dt, \\ A_3(\alpha,\lambda,t,\varphi) &= \int_0^1 \left|t\left(\lambda - t^\alpha\right)\right| (1-t)\,\varphi(1-t)\,dt. \end{aligned}$$

Proof. By using Lemma 1, the power mean inequality, then we get

(2.2)

$$\begin{aligned} &|S_f(x,\lambda,\alpha,t,\varphi;a,b)| \\ &\leq \tfrac{(x-a)^{\alpha+2}}{b-a}\left(\int_0^1 \left|t\left(\lambda - t^\alpha\right)\right| dt\right)^{1-\frac{1}{q}}\left(\int_0^1 \left|t\left(\lambda - t^\alpha\right)\right| \left|f''\left(tx+(1-t)a\right)\right|^q dt\right)^{\frac{1}{q}} \\ &\quad + \tfrac{(b-x)^{\alpha+2}}{b-a}\left(\int_0^1 \left|t\left(\lambda - t^\alpha\right)\right| dt\right)^{1-\frac{1}{q}}\left(\int_0^1 \left|t\left(\lambda - t^\alpha\right)\right| \left|f''\left(tx+(1-t)b\right)\right| dt\right)^{\frac{1}{q}} \\ &= A_1^{1-\frac{1}{q}}(\alpha,\lambda)\left[\tfrac{(x-a)^{\alpha+2}}{b-a}\left(\int_0^1 \left|t(\lambda - t^\alpha)\right| \left|f''\left(tx+(1-t)a\right)\right|^q dt\right)^{\frac{1}{q}}\right. \\ &\quad \left. + \tfrac{(b-x)^{\alpha+2}}{b-a}\left(\int_0^1 \left|t(\lambda - t^\alpha)\right| \left|f''\left(tx+(1-t)b\right)\right|^q dt\right)^{\frac{1}{q}}\right], \end{aligned}$$

where

$$A_1(\alpha,\lambda) = \int_0^1 \left|t\left(\lambda - t^\alpha\right)\right| dt = \left(\frac{\alpha\lambda^{1+\frac{2}{\alpha}}+1}{\alpha+2} - \frac{\lambda}{2}\right).$$

Since $|f''|^q$ is $\varphi-$convex on $[a,b]$, we have

$$\begin{aligned} I_1 &= \int_0^1 |t(\lambda - t^\alpha)| \, |f''(tx+(1-t)a)|^q \, dt \\ &\leq \int_0^1 |t(\lambda - t^\alpha)| \left\{ t\varphi(t) \, |f''(x)|^q + (1-t)\varphi(1-t) \, |f''(a)|^q \right\} dt \qquad (2.3) \\ &= A_2(\alpha,\lambda,t,\varphi) \, |f''(x)|^q + A_3(\alpha,\lambda,t,\varphi) \, |f''(a)|^q , \end{aligned}$$

and similarly we can obtain

$$\begin{aligned} I_2 &= \int_0^1 |t(\lambda - t^\alpha)| \, |f''(tx+(1-t)b)|^q \, dt \\ &\leq \int_0^1 |t(\lambda - t^\alpha)| \left\{ t\varphi(t) \, |f''(x)|^q + (1-t)\varphi(1-t) \, |f''(b)|^q \right\} dt \qquad (2.4) \\ &= A_2(\alpha,\lambda,t,\varphi) \, |f''(x)|^q + A_3(\alpha,\lambda,t,\varphi) \, |f''(b)|^q . \end{aligned}$$

By substituting (2.3) and (2.4) in (2.2), we get

$$\begin{aligned} &|S_f(x,\lambda,\alpha,t,\varphi;a,b)| \\ &\leq \left(\tfrac{\alpha\lambda^{1+\frac{2}{\alpha}}+1}{\alpha+2} - \tfrac{\lambda}{2} \right)^{1-\frac{1}{q}} \left[\tfrac{(x-a)^{\alpha+2}}{b-a} \left\{ |f''(x)|^q \int_0^1 |t(\lambda - t^\alpha)| \, t\varphi(t) \, dt \right. \right. \\ &\left. + |f''(a)|^q \int_0^1 |t(\lambda - t^\alpha)| \, (1-t)\varphi(1-t) \, dt \right\}^{\frac{1}{q}} \\ &+ \tfrac{(b-x)^{\alpha+2}}{b-a} \left\{ |f''(x)|^q \int_0^1 |t(\lambda - t^\alpha)| \, t\varphi(t) \, dt \right. \\ &\left. \left. + |f''(b)|^q \int_0^1 |t(\lambda - t^\alpha)| \, (1-t)\varphi(1-t) \, dt \right\}^{\frac{1}{q}} \right]. \end{aligned}$$

Thus the proof is complated. □

Corollary 1. *Let $\varphi(t) = 1$ in Theorem 1, then we get the following inequality:*

$$\begin{aligned} &|S_f(x,\lambda,\alpha;a,b)| \\ &\leq \left(\tfrac{\alpha\lambda^{1+\frac{2}{\alpha}}+1}{\alpha+2} - \tfrac{\lambda}{2} \right)^{1-\frac{1}{q}} \left[\tfrac{(x-a)^{\alpha+2}}{b-a} \left\{ A_2(\alpha,\lambda) \, |f''(x)|^q + A_3(\alpha,\lambda) \, |f''(a)|^q \right\} \right. \\ &\left. + \tfrac{(b-x)^{\alpha+2}}{b-a} \left\{ A_2(\alpha,\lambda) \, |f''(x)|^q + A_3(\alpha,\lambda) \, |f''(b)|^q \right\} \right]. \end{aligned}$$

Where

$$A_2(\alpha,\lambda) = \int_0^1 |t(\lambda - t^\alpha)| \, tdt = \frac{3-(\alpha+3)\lambda + 2\alpha\lambda^{1+\frac{3}{\alpha}}}{3(\alpha+3)}$$

and

$$\begin{aligned} A_3(\alpha,\lambda) &= \int_0^1 |t(\lambda - t^\alpha)| \, (1-t) \, dt \\ &= \frac{\alpha\lambda^{1+\frac{2}{\alpha}}}{\alpha+2} - \frac{2\lambda^{1+\frac{3}{\alpha}}}{3(\alpha+3)} + \frac{\alpha\lambda}{6} - \frac{\alpha}{(\alpha+2)(\alpha+3)}. \end{aligned}$$

Corollary 2. *If we choose $\varphi(t)=1$ and $x=\frac{a+b}{2}$ in Theorem 1, we can obtain the corollary 2.2, 2.3, 2.4 in ([14]), respectively for $\lambda=\frac{1}{3}$, $\lambda=0$, $\lambda=1$.*

Corollary 3. *Let $\varphi(t)=t^{s-1}$ in Theorem 1, then we have*

$$
\begin{aligned}
&|S_f(x,\lambda,\alpha,t,\varphi;a,b)| \\
&\leq \left(\frac{\alpha\lambda^{1+\frac{2}{\alpha}}+1}{\alpha+2}-\frac{\lambda}{2}\right)^{1-\frac{1}{q}}\left[\frac{(x-a)^{\alpha+2}}{b-a}\left\{|f''(x)|^q A_4(\alpha,\lambda,s)+|f''(a)|^q A_5(\alpha,\lambda,t,\varphi)\right\}^{\frac{1}{q}}\right. \\
&\left.+\frac{(b-x)^{\alpha+2}}{b-a}\left\{|f''(x)|^q A_4(\alpha,\lambda,s)+|f''(b)|^q A_5(\alpha,\lambda,t,\varphi)\right\}^{\frac{1}{q}}\right].
\end{aligned}
$$

Where

$$
\begin{aligned}
A_4(\alpha,\lambda,s) &= 2\frac{\lambda^{\frac{s+2}{\alpha}+1}}{s+2}-2\frac{\lambda^{\frac{s+2}{\alpha}+1}}{\alpha+s+2}+\frac{1}{\alpha+s+2} \\
A_5(\alpha,\lambda,t,\varphi) &= \lambda\beta\left(\lambda^{\frac{1}{\alpha}},2,s+1\right)-\beta\left(\lambda^{\frac{1}{\alpha}},\alpha+2,s+1\right) \\
&\quad+\beta\left(1-\lambda^{\frac{1}{\alpha}},\alpha+2,s+1\right)-\lambda\beta\left(1-\lambda^{\frac{1}{\alpha}},2,s+1\right).
\end{aligned}
$$

Theorem 2. *Let $\varphi:(0,1)\to(0,\infty)$ be a measurable function. For $f:I\subset[0,\infty)\to\mathbb{R}$ be a twice differentiable function on the interior I^0 assume also that $f''\in L_1[a,b]$, where $a,b\in I^0$ with $a<b$. If $|f''|^q$ is $\varphi-$convex on $[a,b]$ for some fixed $q>1$ with $\frac{1}{p}+\frac{1}{q}=1$, then for any $x\in[a,b]$, $\lambda\in[0,1]$ and $\alpha>0$ the following inequality holds*

$$
\begin{aligned}
&|S_f(x,\lambda,\alpha,t,\varphi;a,b)| \\
&\leq B^{\frac{1}{p}}(\alpha,\lambda,p)\left[\frac{(x-a)^{\alpha+2}}{b-a}\left\{\left(|f''(x)|^q+|f''(a)|^q\right)\int_0^1 t\varphi(t)\,dt\right\}^{\frac{1}{q}}\right. \\
&\left.+\frac{(b-x)^{\alpha+2}}{b-a}\left\{\left(|f''(x)|^q+|f''(b)|^q\right)\int_0^1 t\varphi(t)\,dt\right\}^{\frac{1}{q}}\right],
\end{aligned}
\tag{2.5}
$$

where

$$
\begin{aligned}
B(\alpha,\lambda,p) = \frac{\lambda^{\frac{1+p+\alpha p}{\alpha}}}{\alpha}&\left\{\Gamma(1+p)\,\Gamma\left(\tfrac{1+p+\alpha}{\alpha}\right)\ \left({}_2F_1\left(1,1+p,2+p+\tfrac{1+p}{\alpha},1\right)\right)\right. \\
&\left.+\beta\left(1+p,-\tfrac{1+p+\alpha p}{\alpha}\right)-\beta\left(\lambda,1+p,-\tfrac{1+p+\alpha p}{\alpha}\right)\right\},
\end{aligned}
$$

also, for $0<b<c$ and $|z|<1$, ${}_2F_1$ is hypergeometric function defined by

$$
{}_2F_1(a,b,c,z)=\frac{1}{\beta(b,c-b)}\int_0^1 t^{b-1}(1-t)^{c-b-1}(1-zt)^{-a}\,dt.
$$

Proof. By using Lemma 1 and the Hölder inequality, we have the below inequality

$$\begin{aligned}
&|S_f(x,\lambda,\alpha,t,\varphi;a,b)| \\
&\leq \tfrac{(x-a)^{\alpha+2}}{b-a}\left(\int_0^1 |t(\lambda-t^\alpha)|^p\,dt\right)^{\frac{1}{p}}\left(\int_0^1 |f''(tx+(1-t)a)|^q\,dt\right)^{\frac{1}{q}} \\
&+\tfrac{(b-x)^{\alpha+2}}{b-a}\left(\int_0^1 |t(\lambda-t^\alpha)|^p\,dt\right)^{\frac{1}{p}}\left(\int_0^1 |f''(tx+(1-t)b)|^q\,dt\right)^{\frac{1}{q}} \\
&=\left(\int_0^1 |t(\lambda-t^\alpha)|^p\right)^{\frac{1}{p}}\left[\tfrac{(x-a)^{\alpha+2}}{b-a}\left(\int_0^1 |f''(tx+(1-t)a)|^q\,dt\right)^{\frac{1}{q}}\right. \\
&\left.+\tfrac{(b-x)^{\alpha+2}}{b-a}\left(\int_0^1 |f''(tx+(1-t)b)|^q\,dt\right)^{\frac{1}{q}}\right].
\end{aligned} \tag{2.6}$$

Since $|f''|$ is $\varphi-$convex on $[a,b]$, we have

$$\begin{aligned}
\int_0^1 |f''(tx+(1-t)a)|^q\,dt &\leq \int_0^1 t\varphi(t)\,|f''(x)|^q\,dt \\
&+\int_0^1 (1-t)\,\varphi(1-t)\,|f''(a)|^q\,dt \\
&=\left(|f''(x)|^q+|f''(a)|^q\right)\int_0^1 t\varphi(t)\,dt,
\end{aligned} \tag{2.7}$$

and using same tecnique, we get

$$\begin{aligned}
\int_0^1 |f''(tx+(1-t)b)|^q\,dt &\leq \int_0^1 t\varphi(t)\,|f''(x)|^q\,dt \\
&+\int_0^1 (1-t)\,\varphi(1-t)\,|f''(b)|^q\,dt \\
&=\left(|f''(x)|^q+|f''(b)|^q\right)\int_0^1 t\varphi(t)\,dt.
\end{aligned} \tag{2.8}$$

On the other hand we can obtain the following equality;

$$\begin{aligned}
B(\alpha,\lambda,p) &= \int_0^1 |t(\lambda-t^\alpha)|^p\,dt \\
&= \int_0^{\lambda^{\frac{1}{\alpha}}} \{t(\lambda-t^\alpha)\}^p\,dt+\int_{\lambda^{\frac{1}{\alpha}}}^1 \{t(t^\alpha-\lambda)\}^p\,dt \\
&= C_1(\alpha,\lambda,p)+C_2(\alpha,\lambda,p).
\end{aligned} \tag{2.9}$$

By letting $\lambda - t^{\alpha} = u$ and $t^{\alpha} = u$, respectively, we have

(2.10)
$$\begin{aligned} C_1(\alpha,\lambda,p) &= \int_0^{\lambda^{\frac{1}{\alpha}}} \{t(\lambda - t^{\alpha})\}^p \, dt \\ &= \tfrac{1}{\alpha} \int_0^{\lambda} u^p (\lambda - u)^{\frac{1+p-\alpha}{\alpha}} \, du \\ &= \tfrac{1}{\alpha} \int_0^1 \lambda^p y^p \lambda^{\frac{1+p-\alpha}{\alpha}} (1-y)^{\frac{1-\alpha+p}{\alpha}} \lambda dy \\ &= \frac{\lambda^{\frac{p\alpha+1+p}{\alpha}}}{\alpha} \int_0^1 y^p (1-y)^{\frac{1+p}{\alpha}} (1-y)^{-1} \, dy \\ &= \tfrac{\lambda^{\frac{1+p+\alpha p}{\alpha}}}{\alpha} \Gamma(1+p) \Gamma\left(\tfrac{1+p+\alpha}{\alpha}\right)_2 F_1\left(1, 1+p, 2+p+\tfrac{1+p}{\alpha}, 1\right), \end{aligned}$$

and

$$\begin{aligned} C_2(\alpha,\lambda,p) &= \int_{\lambda^{\frac{1}{\alpha}}}^1 \{t(t^{\alpha} - \lambda)\}^p \, dt \\ &= \tfrac{1}{\alpha} \int_{\lambda^u}^1 \tfrac{1+p-\alpha}{\alpha} (u-\lambda)^p \, du \\ &= \tfrac{\lambda^{\frac{1+p+\alpha p}{\alpha}}}{\alpha} \left\{\beta\left(1+p, -\tfrac{1+p+\alpha p}{\alpha}\right) - \beta\left(\lambda, 1+p, -\tfrac{1+p+\alpha p}{\alpha}\right)\right\}. \end{aligned} \tag{2.11}$$

Thus, we get the desired result. □

Corollary 4. *Let $\varphi(t) = 1$ in Theorem 2, then we get the following inequality for any $x \in [a,b], \lambda \in [0,1]$ and $\alpha > 0$;*

$$\begin{aligned} &|S_f(x,\lambda,\alpha,t,\varphi;a,b)| \\ &\leq \left(\int_0^1 |t(\lambda - t^{\alpha})|^p \, dt\right)^{\frac{1}{p}} \left[\tfrac{(x-a)^{\alpha+2}}{b-a} \left\{\tfrac{\left(|f''(x)|^q + |f''(a)|^q\right)}{2}\right\}^{\frac{1}{q}}\right. \\ &\left. + \tfrac{(b-x)^{\alpha+2}}{b-a} \left\{\tfrac{\left(|f''(x)|^q + |f''(b)|^q\right)}{2}\right\}^{\frac{1}{q}}\right]. \end{aligned}$$

Corollary 5. *If we choose $\varphi(t) = 1$ and $x = \frac{a+b}{2}$ in Theorem 2, we can obtain the corollary 2.6, 2.7, 2.8 in ([14]), respectively for $\lambda = \frac{1}{3}$, $\lambda = 0$, $\lambda = 1$.*

Corollary 6. *Let in $\varphi(t) = t^{s-1}$ Theorem 2, then we obtain*

$$\begin{aligned} &|S_f(x,\lambda,\alpha,t,\varphi;a,b)| \\ &\leq \left(\int_0^1 |t(\lambda - t^{\alpha})|^p \, dt\right)^{\frac{1}{p}} \left[\tfrac{(x-a)^{\alpha+2}}{b-a} \left\{\tfrac{\left(|f''(x)|^q + |f''(a)|^q\right)}{s+1}\right\}^{\frac{1}{q}}\right. \\ &\left. + \tfrac{(b-x)^{\alpha+2}}{b-a} \left\{\tfrac{\left(|f''(x)|^q + |f''(b)|^q\right)}{s+1}\right\}^{\frac{1}{q}}\right]. \end{aligned}$$

References

[1] E. F. Beckenbach, Convex functions, *Bull. Amer. Math. Soc.*, **54** (1948) 439-460. http://dx.doi.org/10.1090/s0002-9904-1948-08994-7

[2] Z. Dahmani, On Minkowski and Hermite-Hadamard integral inequalities via fractional integration, *Ann. Funct. Anal.*, 1(**1**) (2010) 51-58. http://dx.doi.org/10.15352/afa/1399900993

[3] S. S. Dragomir, Inequalities of Jensen type for φ-convex functions, *Fasc. Math.* **55** (2015) 35-52

[4] H. Hudzik and L. Maligranda, Some remarks on s-convex functions, Aequationes Math., 48 (1994), no. 1, 100-111.

[5] I. Işcan, K. Bekar, S. Numan, Hermite-Hadamard an Simpson type inequalities for differentiable quasi-geometrically convex func- tions, *Turkish J. of Anal. and Number Theory*, 2(**2**) (2014) 42-46. http://dx.doi.org/10.12691/tjant-2-2-3

[6] I. Işcan, New estimates on generalization of some integral inequalities for ds-convex functions and their applications, *Int. J. Pure Appl. Math.*, 86(4) (2013) 727-746. http://dx.doi.org/10.12732/ijpam.v86i4.11

[7] I. Işcan, Generalization of different type integral inequalities via fractional integrals for functions whose second derivatives absolute value are quasi-convex *Konuralp Journal of Mathematics*, 1(2) (2013) 67-79.

[8] I. Işcan, On generalization of different type integral inequalities for s-convex functions via fractional integrals presented

[9] H. Kavurmaci, M. Avci, M. E. Özdemir, New inequalities of Hermite- Hadamard's type for convex functions with applications, *Journ. of In-equal. and Appl.*, 2011:86 (2011). http://dx.doi.org/10.1186/1029-242x-2011-86

[10] V. G. Mihesan, A generalization of the convexity, *Seminar on Functional Equations, Approx. and Convex, Cluj-Napoca, Romania* (1993).

[11] M. E. Özdemir, M. Avic, H. Kavurmaci, Hermite-Hadamard type inequalities for s-convex and s-concave functions via fractional integrals, arXiv:1202.0380v1[math.CA].

[12] J. Park, Generalization of some Simpson-like type inequalities via differentiable s-convex mappings in the second sense, *In- ter. J. of Math. and Math. Sci.*, 2011 Art No: 493531, 13 pages. http://dx.doi.org/10.1155/2011/493531

[13] Jaekeun Park, Some new Hermite-Hadamard-like type inequalities on geometrically convex functions, *Inter. J. of Math. Anal.*, 8(**16**) (2014),793-802. http://dx.doi.org/10.12988/ijma.2014.4243

[14] Jaekeun Park, On Some Integral Inequalities for Twice Differentiable Quasi-Convex and Convex Functions via Fractional Integrals, *Applied Mathematical Sciences*, Vol. 9(62) (2015), 3057-3069 HIKARI Ltd, www.m-hikari.com. http://dx.doi.org/10.12988/ams.2015.53248.

[15] Samko, S.G., Kilbas A.A. and Marichev, O.I., Fractional Integrals and Derivatives, Theory and Applications, *Gordon and Breach*, 1993, ISBN 2881248640.

[16] M. Z. Sarikaya, H. Ogunmez, On new inequalities via Riemann-Liouville fractional integration, *Abstract and applied analysis*, 2012 (2012) 10 pages, Art ID:428983. http://dx.doi.org/10.1155/2012/428983

[17] M. Z. Sarikaya, E. Set, H. Yildiz, N. Basak, Hermite- Hadamard's inequalities for fractional integrals and related frac- tional inequalities, *Math. and Comput. Model.*, 2011 (2011). http://dx.doi.org/10.1016/j.mcm.2011.12.048

[18] E. Set, M. Z. Sarikaya, M. E. Özdemir, Some Ostrowski's type Inequalities for functions whose second derivatives are s-convex in the second sense, arXiv:1*006.24* 88v1 [*math.* CA] 12 *June 2010.*

[19] E. Set, E. Ozdemir, M. Z. Sarikaya, F. Karako, Hermite-Hadamard type inequalities for mappings whose derivatives are s-convex in the second sense via fractional integrals, *Khayyam J. Math.,* 1(1) (2015) 62-70.

[20] Gh. Toader, On a generalization of the convexity, *Mathematica*, 30(53) (1988), 83-87.

[21] M.Tunc, On some new inequalities for convex functions, *Turk. J. Math.*, **35** (2011), 1-7.

[22] M. Tunc, H. Yildirim, *On MT-Convexity, arXiv: 1205.5453 [math. CA] 24 May 2012*

[Department of Mathematics, Faculty of Science, University of Cumhuriyet, 58140, Sivas, Turkey
E-mail address: mesra@cumhuriyet.edu.tr

[Department of Mathematics, Faculty of Science and Arts, University of Kahramanmaraş Sütçü İmam, 46100, Kahramanmaraş, Turkey
E-mail address: abdullahmat@gmail.com

[Department of Mathematics, Faculty of Science and Arts, University of Kahramanmaraş Sütçü İmam, 46100, Kahramanmaraş, Turkey
E-mail address: hyildir@ksu.edu.tr